Your Guide to Understanding Genetic Conditions

GLDC gene

glycine decarboxylase

Normal Function

The *GLDC* gene provides instructions for making an enzyme called glycine dehydrogenase. This enzyme is one of four components (subunits) that make up a large complex called glycine cleavage enzyme. Within cells, this complex is active in specialized energy-producing centers called mitochondria.

As its name suggests, glycine cleavage enzyme processes a molecule called glycine by cutting (cleaving) it into smaller pieces. Glycine is an amino acid, which is a building block of proteins. This molecule also acts as a neurotransmitter, which is a chemical messenger that transmits signals in the brain. The breakdown of excess glycine is necessary for the normal development and function of nerve cells in the brain and spinal cord.

Health Conditions Related to Genetic Changes

glycine encephalopathy


Mutations in the *GLDC* gene account for about 80 percent of all cases of glycine encephalopathy. More than 40 mutations have been identified in affected individuals. Many of these genetic changes alter single amino acids in glycine dehydrogenase. For example, the most common *GLDC* mutation in the Finnish population replaces the amino acid serine with the amino acid isoleucine at position 564 in the enzyme (also written as Ser564lle or S564l). Other mutations insert or delete genetic material in the *GLDC* gene, or disrupt how genetic information from the gene is spliced together to make a blueprint for producing glycine dehydrogenase.

Some *GLDC* mutations lead to the production of a nonfunctional version of glycine dehydrogenase, while other mutations reduce but do not eliminate the enzyme's activity. When an altered version of this enzyme is incorporated into the glycine cleavage enzyme complex, it prevents the complex from breaking down glycine properly. As a result, excess glycine can build up to toxic levels in the body's organs and tissues. Damage caused by harmful amounts of this molecule in the brain and spinal cord is responsible for the intellectual disability, seizures, and breathing difficulties characteristic of glycine encephalopathy.

Chromosomal Location

Cytogenetic Location: 9p24.1, which is the short (p) arm of chromosome 9 at position 24.1

Molecular Location: base pairs 6,532,464 to 6,645,692 on chromosome 9 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- GCE
- GCSP
- GCSP_HUMAN
- glycine cleavage system protein P
- glycine decarboxylase P-protein
- glycine dehydrogenase (decarboxylating)
- glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)
- HYGN1
- NKH

Additional Information & Resources

Educational Resources

- Basic Neurochemistry (sixth edition, 1999): Nonketotic hyperglycinemia is caused by deficiencies in the glycine-cleavage system https://www.ncbi.nlm.nih.gov/books/NBK27969/
- Nomenclature Committee of the International Union of Biochemistry and Molecular Biology: Glycine Cleavage System http://www.chem.gmul.ac.uk/iubmb/enzyme/reaction/AminoAcid/GlyCleave.html

GeneReviews

 Glycine Encephalopathy https://www.ncbi.nlm.nih.gov/books/NBK1357

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28GLDC%5BTIAB%5D%29+OR+%28glycine+dehydrogenase%5BTIAB%5D%29%29+OR+%28glycine+decarboxylase%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D

OMIM

 GLYCINE DECARBOXYLASE http://omim.org/entry/238300

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology http://atlasgeneticsoncology.org/Genes/GC_GLDC.html
- ClinVar https://www.ncbi.nlm.nih.gov/clinvar?term=GLDC%5Bgene%5D
- HGNC Gene Symbol Report http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/ hgnc_data.php&hgnc_id=4313
- NCBI Gene https://www.ncbi.nlm.nih.gov/gene/2731
- UniProt http://www.uniprot.org/uniprot/P23378

Sources for This Summary

- Applegarth DA, Toone JR. Glycine encephalopathy (nonketotic hyperglycinemia): comments and speculations. Am J Med Genet A. 2006 Jan 15;140(2):186-8.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/16353254
- Conter C, Rolland MO, Cheillan D, Bonnet V, Maire I, Froissart R. Genetic heterogeneity of the GLDC gene in 28 unrelated patients with glycine encephalopathy. J Inherit Metab Dis. 2006 Feb; 29(1):135-42.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/16601880
- Dinopoulos A, Matsubara Y, Kure S. Atypical variants of nonketotic hyperglycinemia. Mol Genet Metab. 2005 Sep-Oct;86(1-2):61-9. Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/16157495
- GeneReview: Glycine Encephalopathy https://www.ncbi.nlm.nih.gov/books/NBK1357

- Kure S, Kato K, Dinopoulos A, Gail C, DeGrauw TJ, Christodoulou J, Bzduch V, Kalmanchey R, Fekete G, Trojovsky A, Plecko B, Breningstall G, Tohyama J, Aoki Y, Matsubara Y. Comprehensive mutation analysis of GLDC, AMT, and GCSH in nonketotic hyperglycinemia. Hum Mutat. 2006 Apr; 27(4):343-52.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/16450403
- Nakai T, Nakagawa N, Maoka N, Masui R, Kuramitsu S, Kamiya N. Structure of P-protein of the glycine cleavage system: implications for nonketotic hyperglycinemia. EMBO J. 2005 Apr 20;24(8): 1523-36. Epub 2005 Mar 24.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/15791207
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1142568/
- Sellner L, Edkins E, Greed L, Lewis B. Detection of mutations in the glycine decarboxylase gene in patients with nonketotic hyperglycinaemia. Mol Genet Metab. 2005 Feb;84(2):167-71. Epub 2004 Nov 23.

Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/15670722

Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/gene/GLDC

Reviewed: April 2007 Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services